

4.163 Global scale variability of the mineral dust shortwave and longwave refractive index: a new dataset of in situ measurements for climate modelling and remote sensing.

Presenting Author:

Claudia Di Biagio, LISA, CNRS, UPEC and UPD, IPSL, Créteil, France,
claudia.dibiagio@lisa.u-pec.fr

Co-Authors:

Paola Formenti, LISA, CNRS, UPEC and UPD, IPSL, Créteil, France
Yves Balkanski, LSCE, CEA CNRS UVSQ, 91191, Gif sur Yvette, France
Lorenzo Caponi, LISA, CNRS, UPEC and UPD, IPSL, Créteil, France
Mathieu Cazaunau, LISA, CNRS, UPEC and UPD, IPSL, Créteil, France
Edouard Pangui, LISA, CNRS, UPEC and UPD, IPSL, Créteil, France
Emilie Journet, LISA, CNRS, UPEC and UPD, IPSL, Créteil, France
Sophie Nowak, Plateforme RX UFR de chimie, UPD, Paris, France
Sandrine Caquineau, IRD-Sorbonne Universités (UPMC) – CNRS-MNHN, LOCEAN, IRD France-Nord, Bondy, France
Meinrat Andreae, Max Planck Institute for Chemistry, Mainz, Germany
Konrad Kandler, Technische Universität Darmstadt, Darmstadt, Germany
Thuraya Saeed, College of Basic Education, PAAET, Al-Ardeya, Kuwait
Stuart Piketh, North-West University, Potchefstroom, South Africa
David Seibert, Walden University, Minneapolis, Minnesota, USA
Earle Williams, Parsons Laboratory, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA
Jean-Francois Doussin, LISA, CNRS, UPEC and UPD, IPSL, Créteil, France

Abstract:

Modeling the interaction of dust with short-wave (SW) and long-wave (LW) radiation is still a challenge because of the scarcity of information on the complex refractive index ($n-ik$) of dust and its variability as a function of the particle mineralogical composition, which depends on the specific emission source. As a consequence, to date, climate models and remote sensing retrievals use a spatially invariant value for the dust SW and LW refractive index.

In this study, the global variability of the mineral dust SW (0.37-0.95 μm) and LW (2-16 μm) refractive index as a function of its mineralogical composition is explored by in situ measurements in a large smog chamber. Investigated dust aerosol samples were issued from major desert sources worldwide, including the African Sahara and Sahel, Eastern Asia, the Middle East, Southern Africa, Australia, and the Americas.

Results from the present study provide a regional mapping of the SW and LW optical properties by dust and show that the imaginary part of the refractive index largely varies for the different source areas due to the change in the particle composition. In the LW k varies between ~ 0.001 and 0.92 due to changes in the clays, quartz, and calcite content in dust. In the SW range k varies in the range $\sim 0.002-0.030$ at $0.37 \mu\text{m}$ and $0.0005-0.005$

at 0.95 μm in relation to changes of the iron dust content.

A linear relationship between the magnitude of k at specific wavelengths and the mass concentration of minerals absorbing at these wavelengths was found. Predictive rules could be thus established to estimate the SW and LW refractive index of dust in specific bands based on an assumed or predicted mineralogical composition. These rules can be used to implement regional-dependent SW and LW refractive indices in models and remote sensing retrieval algorithms.